

GRRM NEWS No. 004

GRRM NEWS reports recent developments of the ADD/SHS algorithm and its applications. The newest version of the GRRM program is **GRRM1.22**, which was released on 4 January 2010, and its minor change has been made on 3 May 2010.

(1) The major improvement in **GRRM1.22** is an extension of disposable atoms up to Rg (atomic number=111). Many metal atoms can now be handled by the GRRM program.

(2) In connection with the above improvement, partial coordinates of more than two atoms in the system can be frozen as “a field” of the remaining part.

This “field” can easily be introduced by adding series of atomic coordinates at the end of the coordinates for the flexible part of the system to be optimized by placing just one line starting from a word of “FIELD” as a separator between the flexible part and the frozen part.

Before applying the GRRM option, the MIN option (geometry optimization) should be used to obtain optimized structures as the starting geometries for the GRRM option.

With using this technique, a large catalytic system with BINAP could be handled, and further applications to surface reactions are under progress.

(3) A graphical display (GDSP) system for GRRM calculations has been developed. GRRM-GDSP is a data handling system, which automatically creates a system of web pages for viewing lists of obtained geometries and three-dimensional structures based on output data from the GRRM program.

Animations around transition structures along IRC pathways can also be produced automatically.

(4) An general extension of the scaled hyper sphere search (SHS) algorithm to the more general problems has been under progress. It has been proved that the SHS algorithm can be used for global optimization for a smooth system in which adjacent minima have mutual influences with a perceivable distinction between them.

GRRM1.22 is an *updated* version of the first program based on the SHS algorithm for an automated exploration of reaction pathways by using energies from solutions of $H\Psi = E\Psi$.

GRRM 1.22 copes with various problems in chemistry by automated exploration of reaction pathways.

- **GRRM 1.22 automatically explores unknown isomers.**
- **GRRM 1.22 automatically explores unknown synthetic routes.**
- **GRRM 1.22 automatically explores unknown dissociation channels.**

GRRM 1.22 develops an unexplored world of chemistry by elucidating unknown reaction networks.

- **GRRM 1.22 is useful for production of the Atlas for the chemical world.**
- **GRRM 1.22 is useful for design of new chemical compounds and reactions.**
- **GRRM 1.22 is useful for designing new tactics for energy/environment problems.**
- **GRRM 1.20 is useful for elucidation of catalysis and design of new catalysts.**

GRRM 1.22 is an epoch-making program of potential analyses for the following problems.

- **Normal coordinate analysis** Normal coordinate calculations can be made at arbitrary structures. Optionally, enthalpy and Gibbs energies can also be obtained.
- **Optimization of equilibrium structures** Equilibrium structures can be optimized by SIRFO and BFGS methods.
- **Optimization of transition structures** Transition structures can be optimized by SIRFO and Bofill's methods.
- **IRC search** IRC can be traced by Page and McIver methods.
- **GRRM search** Global reaction route mapping (**GRRM**) can be made for the potential surface of a given chemical formula. Starting from an equilibrium structure, automated search of dissociation and isomerization can be performed to explore **GRRM** corresponding to the Atlas of chemical reaction routes. Optionally, exploration of reaction routes can be made for the limited region around a particular structure.
- **One step TS search** An efficient search of the reaction pathway via a transition structure (TS) between a reactant and a product can be made automatically without initial guess, and this technique is much more rapid and applicable than any other methods.
- **Intermediate search** Intermediates between a pair of isomers can be found, even if they are far apart. The SHS method in the hypersphere-contraction-mode enables us to explore multi-step reaction pathways, even if they amount to several tens of steps.
- **Large ADD following (LADD)** A very efficient search of lower lying structures can be made by the LADD algorithm, which is especially suitable for systems with huge numbers of isomers.
- **ONIOM and various QM&MM methods** ONIOM as well as various methods available in the Gaussian program can be used as options in combination with the above techniques.

Program Package & Requirement for GRRM 1.22


GRRM 1.22 utilizes energies obtained by **Gaussian03** (Gaussian09 can also be used directly).

Conventional packages other than Gaussian03 can also be used with additional data handling.

GRRM 1.22 can be used under a Linux/Unix environment.

GRRM 1.22 can be used for research and education, after application to the following address by

E-mail. ohnok@m.tains.tohoku.ac.jp



SHS-GRRM Publication List

- 1) A Scaled Hypersphere Search Method for the Topography of Reaction Pathways on the Potential Energy Surface, K. Ohno and S. Maeda, *Chem. Phys. Lett.* 384(4-6), 277-282 (2004).
- 2) Ab initio Studies on Synthetic Routes of Glycine from Simple Molecules via Ammonolysis of Acetolactone: Applications of the Scaled Hypersphere Search Method, S. Maeda and K. Ohno, *Chemistry Letters* 33, 1372-1373 (2004).
- 3) No Activation Barrier Synthetic Route of Glycine from Simple Molecules (NH₃, CH₂, and CO₂) via Carboxylation of Ammonium Ylide: a Theoretical Study by the Scaled Hypersphere Search Method, S. Maeda and K. Ohno, *Chem. Phys. Lett.* 398 (1-3), 240-244 (2004).
- 4) A New Approach for Finding a Transition State Connecting a Reactant and a Product without Initial Guess: Applications of the Scaled Hypersphere Search method to Isomerization Reactions of HCN, (H₂O)₂, and Alanine Dipeptide, S. Maeda and K. Ohno, *Chem. Phys. Lett.* 404(1-3), 95-99 (2005).
- 5) Global Mapping of Equilibrium and Transition Structures on Potential Energy Surfaces by the Scaled Hypersphere Search Method: Application to ab initio Surfaces of Formaldehyde and Propyne Molecules, S. Maeda and K. Ohno, *J. Phys. Chem. A* 109(25), 5742-5753 (2005).
- 6) Global Investigation on Potential Energy Surface of CH₃CN: Application of the Scaled Hypersphere Search Method, Xia Yang, Satoshi Maeda, and Koichi Ohno, *J. Phys. Chem. A* 109(32), 7319-7328 (2005).
- 7) A Scaled Hypersphere Interpolation Technique for Efficient Construction of Multidimensional Potential Energy Surfaces, S. Maeda, Y. Watanabe, and K. Ohno, *Chem. Phys. Lett.* 414(4-6), 265-270 (2005).
- 8) Global Analysis of Reaction Pathways on the Potential Energy Surface of Cyanoacetylene by the Scaled Hypersphere Search Method, X. Yang, S. Maeda, and K. Ohno, *Chem. Phys. Lett.* 418(1-3), 208-216 (2006).
- 9) Generation Mechanisms of Amino Acids in the Interstellar Space via Reactions between Closed-Shell Molecules, S. Maeda and K. Ohno, *Astrophys. J.* 640, 823 (2006).
- 10) D-L Conversion Pathways between Optical Isomers of Alanine: Application of the Scaled Hypersphere Method to Explore Unknown Reaction Routes in a Chiral System, K. Ohno and S. Maeda, *Chemistry Letters* 35(5), 492-493 (2006).
- 11) Conversion Pathways between a Fullerene and a Ring among C₂₀ Clusters: Remarkable Difference in Local Potential Energy Landscapes around the Fullerene and the Ring, S. Maeda and K. Ohno, *J. Chem. Phys.* 124, 174306-(1,7) (2006).
- 12) Global Reaction Route Mapping on Potential Energy Surfaces of Formaldehyde, Formic Acid, and their Metal Substituted Analogues, K. Ohno and S. Maeda, *J. Phys. Chem. A* 110(28), 8933-8941 (2006).
- 13) Structures of Water Octamers (H₂O)₈: Exploration on Ab Initio Potential Energy Surfaces by the Scaled Hypersphere Search Method, S. Maeda and K. Ohno, *J. Phys. Chem. A* 111(20), 4527-4534 (2007).
- 14) Insight into Global Reaction Mechanism of [C₂, H₄, O] System from ab initio Calculations by the Scaled Hypersphere Search Method, X. Yang, S. Maeda, and K. Ohno, *J. Phys. Chem. A* 111(23), 5099-5110 (2007).

- 15) A Computational Study of Titanocene-Catalyzed Dehydrocoupling of $\text{Me}_2\text{NH-BH}_3$ Adduct: An Intramolecular, Stepwise Mechanism, Y. Luo and K. Ohno, *Organometallics* 26, 3597-3600 (2007).
- 16) Quantum Chemistry Study of $\text{H}^+(\text{H}_2\text{O})_8$: A Global Search for Its Isomers by the Scaled Hypersphere Search Method and Its Thermal Behavior, Y. Luo, S. Maeda, and K. Ohno, *J. Phys. Chem. A* 111(42), 10732-10737 (2007).
- 17) Global Reaction Route Mapping on Potential Energy Surfaces of C_2H_7^+ and C_3H_9^+ , Y. Watanabe, S. Maeda, and K. Ohno, *Chem. Phys. Lett.* 447/1-3, 21-26 (2007).
- 18) Automated Exploration of Absorption Structures of an Organic Molecule on $\text{RuH}_2\text{-BINAP}$ by the ONIOM Method and the Scaled Hypersphere Search Method, S. Maeda, and K. Ohno, *J. Phys. Chem. A* 111, 13168-13171 (2007).
- 19) Microsolvation of Hydrogen Sulfide: Exploration of $\text{H}_2\text{S} \cdot (\text{H}_2\text{O})_n$ and $\text{SH}^- \cdot \text{H}_3\text{O}^+ \cdot (\text{H}_2\text{O})_{n-1}$ ($n=5-7$) Cluster Structures on Ab Initio Potential Energy Surfaces by the Scaled Hypersphere Search Method, S. Maeda and K. Ohno, *J. Phys. Chem. A* 112(13), 2962-2968 (2008).
- 20) Finding Important Anharmonic Terms in the Sixth-Order Potential Energy Function by the Scaled Hypersphere Search Method: An Application to Vibrational Analyses of Molecules and Clusters, S. Maeda, Y. Watanabe, and K. Ohno, *J. Chem. Phys.* 128, 144111-(1,11) (2008).
- 21) DFT Study on Isomerization and Decomposition of Cuprous Dialkyldithiophosphate and Its Reaction with Alkyl Radical, Y. Luo, S. Maeda, and K. Ohno, *J. Phys. Chem. A* 112(25), 5720-5726 (2008).
- 22) A New Global Reaction Route Map on the Potential Energy Surface of H_2CO with Unrestricted Level, S. Maeda and K. Ohno, *Chem. Phys. Lett.* 460, 55-58 (2008).
- 23) Intramolecular Vibrational Frequencies of Water Clusters $(\text{H}_2\text{O})_n$ ($n=2-5$): Anharmonic Analyses Using Accurate Potential Functions based on the Scaled Hypersphere Search Method, Y. Watanabe, S. Maeda, and K. Ohno, *J. Chem. Phys.* 129, 074315-(1,9) (2008).
- 24) Automated Exploration of Reaction Channels, K. Ohno and S. Maeda, *Physica Scripta* 78, 058122 (8pp) (2008).
- 25) Decomposition of Alkyl Hydroperoxide by a Copper (I) Complex: Insights from Density Functional Theory, Y. Luo, S. Maeda, and K. Ohno, *Tetrahedron Lett.* 49, 6841-6845 (2008).
- 26) Lowest Transition State for the Chirality-Determining Step in $\text{Ru}\{(\text{R})\text{-BINAP}\}$ -Catalyzed Asymmetric Hydrogenation of Methyl-3-Oxobutanoate, S. Maeda and K. Ohno, *J. Am. Chem. Soc.* 130(51), 17228-17229 (2008).
- 27) Water-Catalyzed Gas-Phase Reaction of Formic Acid with Hydroxyl Radical: A Computational Investigation, Y. Luo, S. Maeda, and K. Ohno, *Chem. Phys. Lett.* 469(1-3), 57-61 (2009).
- 28) Automated Exploration of Stable Isomers of $\text{H}^+(\text{H}_2\text{O})_n$ ($n=5-7$) via Ab Initio Calculations: An Application of the Anharmonic Downward Distortion Following Algorithm, Y. Luo, S. Maeda, and K. Ohno, *J. Comp. Chem.* 30(6), 952-961 (2009).
- 29) Automated Global Mapping of Minimum Energy Points on Seams of Crossing by the Anharmonic Downward Distortion Following Method: A Case Study on H_2CO , S. Maeda, K. Ohno and K. Morokuma, *J. Phys. Chem. A* 113(8), 1704-1710 (2009).
- 30) Systematic Search for Isomerization Pathways of Hexasilabenzene for Finding its Kinetic Stability, M. Moteki, S. Maeda, and K. Ohno, *Organometallics* 28(7), 2218-2224 (2009).

- 31) An Automated and Systematic Transition-Structure Explorer in Large Flexible Molecular Systems Based on Combined Global Reaction Route Mapping and Microiteration Methods. S. Maeda K. Ohno, and K. Morokuma, *J. Chem. Theory Comput.* **5**, 2734-2743 (2009).
- 32) A Systematic Study on the RuHCl-BINAP Catalyzed Asymmetric Hydrogenation Mechanism by the Global Reaction Route Mapping Method, Koichi Ohno and Satoshi Maeda, *J. Mol. Cat. A Chemical* **324**, 133-140 (2010).
- 33) Synthesis and Structures of Stable Base-Free Dialkylsilanimines, T. Iwamoto, N. Ohnishi, Z. Gui, S. Ishida, H. Isobe, S. Maeda, K. Ohno, and M. Kira, *New J. Chem.* (accepted: in press)
- 34) Updated Branching Plane for Finding Conical Intersections without Coupling Derivative Vectors, S. Maeda, K. Ohno, and K. Morokuma, *J. Chem. Theory Comput.* (accepted: in press)
- 35) A Theoretical Study on the Photodissociation of Acetone: Insight into the Slow Intersystem Crossing and Exploration of Nonadiabatic Pathways to the Ground State, S. Maeda, K. Ohno, and K. Morokuma, *J. Phys. Chem. Letters* (accepted: in press)