

GRRM NEWS No. 006


GRRM NEWS reports recent developments of the ADD/SHS algorithm and its applications. A major revision has been made for the GRRM program. The newest version is **GRRM11.01**, which is now released for academic users.

(1) One of the major improvements in *GRRM11* is parallel treatments of the GRRM procedure around equilibrium (EQ) points. Although the standard GRRM treatments are processed in series from one EQ point to other unprocessed EQ points. Now in *GRRM11*, the SHS search of reaction channels leading to TS or DC around EQ can be made for several EQ points simultaneously in a parallel mode. This is especially suitable for high performance servers with multi-cores, although this new function is limited to only one-node. This parallel version of GRRM considerably reduces computation time for the global mapping of the entire reaction channels.


(2) Microiteration technique is introduced in *GRRM11* to reduce computational demands for large systems. A giant system with more than 300 atoms could be handled by using QM/MM methods [Ref.32].

(3) Excited-state potential surfaces can be studied by *GRRM11*. Minimum points on seams of crossings as well as conical intersections can be searched very effectively, which helps to consider unknown chemistry in photochemical processes [Refs. 30, 35, 36].


(4) *GRRM11* has been strengthened its ability for studying important problems. Surface processes can be treated with fixing geometries of some atoms in the substrate [Ref. 37]. Imposition of special conditions on some bonds limits chemical structures to be searched, and such treatments considerably reduce computation time to obtain interesting reaction channels leading to aimed chemical compounds.



In addition to the new version of *GRRM11*, a graphical display (GDSP) system for GRRM calculations has been developed. GRRM-GDSP is a data handling system, which automatically creates a system of Web pages for viewing lists of obtained geometries and three-dimensional structures based on output data from the GRRM program. Animations around transition structures along IRC pathways can also be produced automatically. Dissociated products are listed for all dissociation channels, and synthetic reaction routes without byproducts can easily be found on the Web representation.



A new GRRM interface has been created. Everything, from input-data preparation and submission of a GRRM-JOB to confirmation of the searched results as well as creation of visualized Web Pages of the GRRM output data, can be handled on a Web Page in the WebGRRM system.



GRRM11 is an *updated* version of the GRRM program based on the SHS algorithm for an automated exploration of reaction pathways by using energies from solutions of $H\Psi = E\Psi$.

GRRM 11 copes with various problems in chemistry by automated exploration of reaction pathways.

- **GRRM 11 automatically explores unknown isomers.**
- **GRRM 11 automatically explores unknown synthetic routes.**
- **GRRM 11 automatically explores unknown dissociation channels.**

GRRM 11 develops an unexplored world of chemistry by elucidating unknown reaction networks.

- **GRRM 11 is useful for production of the Atlas for the chemical world.**
- **GRRM 11 is useful for design of new chemical compounds and reactions.**
- **GRRM 11 is useful for designing new tactics for energy/environment problems.**
- **GRRM 11 is useful for elucidation of catalysis and design of new catalysts.**

GRRM 11 is an epoch-making program of potential analyses for the following problems.

- **Normal coordinate analysis** Normal coordinate calculations can be made at arbitrary structures. Optionally, enthalpy and Gibbs energies can also be obtained.
- **Optimization of equilibrium structures** Equilibrium structures can be optimized by **RFO** and **BFGS** methods.
- **Optimization of transition structures** Transition structures can be optimized by **RFO** and **Bofill's** methods.
- **IRC search** IRC can be traced by **Page** and **McIver** methods.
- **GRRM search** Global reaction route mapping (**GRRM**) can be made for the potential surface of a given chemical formula. Starting from an equilibrium structure, automated search of dissociation and isomerization can be performed to explore **GRRM** corresponding to the Atlas of chemical reaction routes. Optionally, exploration of reaction routes can be made for the limited region around a particular structure.
- **One step TS search** An efficient search of the reaction pathway via a transition structure (**TS**) between a reactant and a product can be made automatically without initial guess, and this technique is much more rapid and applicable than any other methods.
- **Intermediate search** Intermediates between a pair of isomers can be found, even if they are far apart. The **SHS** method in the hypersphere-contraction-mode enables us to explore multi-step reaction pathways, even if they amount to several tens of steps.
- **Large ADD following (LADD)** A very efficient search of lower lying structures can be made by the **LADD** algorithm, which is especially suitable for systems with huge numbers of isomers.
- **ONIOM and various QM&MM methods** **ONIOM** as well as various methods available in the Gaussian program can be used as options in combination with the above techniques.
- **Microiteration** In connection with **QM&MM**, the microiteration technique can be used for drastically reducing computational demands, and it extends the range of **GRRM** considerably.
- **Limited Search** Limited searches fixing geometries of some atoms as well as imposing special conditions on some bonds extend abilities of the **GRRM** method to many important problems.
- **Excited-State analyses** Minimum energy points on seams of crossings can be searched. With a help of **Molpro**, excited potential energy surfaces can be searched more effectively.

Program Package & Requirement for GRRM 11


GRRM 11 utilizes energies obtained by **Gaussian09** (Gaussian03 can also be used directly).

Conventional packages other than Gaussian09/03 can also be used with additional data handling.

GRRM 11 can be used under a Linux/Unix environment.

GRRM 11 can be used for research and education, after application to the following address by E-mail.

ohnok@m.tohoku.ac.jp



SHS-GRRM Publication List

- 1) A Scaled Hypersphere Search Method for the Topography of Reaction Pathways on the Potential Energy Surface.
K. Ohno and S. Maeda, *Chem. Phys. Lett.* 384(4-6), 277-282 (2004).
- 2) Ab initio Studies on Synthetic Routes of Glycine from Simple Molecules via Ammonolysis of Acetolactone: Applications of the Scaled Hypersphere Search Method.
S. Maeda and K. Ohno, *Chemistry Letters* 33, 1372-1373 (2004).
- 3) No Activation Barrier Synthetic Route of Glycine from Simple Molecules (NH₃, CH₂, and CO₂) via Carboxylation of Ammonium Ylide: a Theoretical Study by the Scaled Hypersphere Search Method.
S. Maeda and K. Ohno, *Chem. Phys. Lett.* 398 (1-3), 240-244 (2004).
- 4) A New Approach for Finding a Transition State Connecting a Reactant and a Product without Initial Guess: Applications of the Scaled Hypersphere Search method to Isomerization Reactions of HCN, (H₂O)₂, and Alanine Dipeptide.
S. Maeda and K. Ohno, *Chem. Phys. Lett.* 404(1-3), 95-99 (2005).
- 5) Global Mapping of Equilibrium and Transition Structures on Potential Energy Surfaces by the Scaled Hypersphere Search Method: Application to ab initio Surfaces of Formaldehyde and Propyne Molecules.
S. Maeda and K. Ohno, *J. Phys. Chem. A* 109(25), 5742-5753 (2005).
- 6) Global Investigation on Potential Energy Surface of CH₃CN: Application of the Scaled Hypersphere Search Method.
Xia Yang, Satoshi Maeda, and Koichi Ohno, *J. Phys. Chem. A* 109(32), 7319-7328 (2005).
- 7) A Scaled Hypersphere Interpolation Technique for Efficient Construction of Multidimensional Potential Energy Surfaces.
S. Maeda, Y. Watanabe, and K. Ohno, *Chem. Phys. Lett.* 414(4-6), 265-270 (2005).
- 8) Global Analysis of Reaction Pathways on the Potential Energy Surface of Cyanoacetylene by the Scaled Hypersphere Search Method.
X. Yang, S. Maeda, and K. Ohno, *Chem. Phys. Lett.* 418(1-3), 208-216 (2006).
- 9) Generation Mechanisms of Amino Acids in the Interstellar Space via Reactions between Closed-Shell Species: Significance of Higher Energy Isomers in Molecular Evolution.
S. Maeda and K. Ohno, *Astrophys. J.* 640, 823 (2006).
- 10) D-L Conversion Pathways between Optical Isomers of Alanine: Application of the Scaled Hypersphere Method to Explore Unknown Reaction Routes in a Chiral System.
K. Ohno and S. Maeda, *Chemistry Letters* 35(5), 492-493 (2006).
- 11) Conversion Pathways between a Fullerene and a Ring among C₂₀ Clusters by a Sphere Contracting Walk Method: Remarkable Difference in Local Potential Energy Landscapes around the Fullerene and the Ring.
S. Maeda and K. Ohno, *J. Chem. Phys.* 124, 174306-(1,7) (2006).
- 12) Global Reaction Route Mapping on Potential Energy Surfaces of Formaldehyde, Formic Acid, and their Metal Substituted Analogues.
K. Ohno and S. Maeda, *J. Phys. Chem. A* 110(28), 8933-8941 (2006).
- 13) Global Mapping of Small Carbon Clusters Using the Scaled Hypersphere Search Method.
B. Hajgato, S. Maeda, and K. Ohno, *AIP Conference Proceedings* 855, 296-304 (2006).
- 14) Structures of Water Octamers (H₂O)₈: Exploration on Ab Initio Potential Energy Surfaces by

- the Scaled Hypersphere Search Method.
S. Maeda and K. Ohno, *J. Phys. Chem. A* 111(20), 4527-4534 (2007).
- 15) Insight into Global Reaction Mechanism of $[C_2, H_4, O]$ System from ab initio Calculations by the Scaled Hypersphere Search Method.
X. Yang, S. Maeda, and K. Ohno, *J. Phys. Chem. A* 111(23), 5099-5110 (2007).
- 16) Computational Study of Titanocene-Catalyzed Dehydrocoupling of the Adduct Me_2NH-BH_3 : An Intramolecular, Stepwise Mechanism.
Y. Luo and K. Ohno, *Organometallics* 26, 3597-3600 (2007).
- 17) Quantum Chemistry Study of $H^+(H_2O)_8$: A Global Search for Its Isomers by the Scaled Hypersphere Search Method and Its Thermal Behavior.
Y. Luo, S. Maeda, and K. Ohno, *J. Phys. Chem. A* 111(42), 10732-10737 (2007).
- 18) Global Reaction Route Mapping on Potential Energy Surfaces of $C_2H_7^+$ and $C_3H_9^+$.
Y. Watanabe, S. Maeda, and K. Ohno, *Chem. Phys. Lett.* 447/1-3, 21-26 (2007).
- 19) Automated Exploration of Absorption Structures of an Organic Molecule on RuH_2 -BINAP by the ONIOM Method and the Scaled Hypersphere Search Method
S. Maeda, and K. Ohno, *J. Phys. Chem. A* 111, 13168-13171 (2007).
- 20) Microsolvation of Hydrogen Sulfide: Exploration of $H_2S \cdot (H_2O)_n$ and $SH^- \cdot H_3O^+ \cdot (H_2O)_{n-1}$ ($n=5-7$) Cluster Structures on Ab Initio Potential Energy Surfaces by the Scaled Hypersphere Search Method.
S. Maeda and K. Ohno, *J. Phys. Chem. A* 112(13), 2962-2968 (2008).
- 21) Finding Important Anharmonic Terms in the Sixth-Order Potential Energy Function by the Scaled Hypersphere Search Method: An Application to Vibrational Analyses of Molecules and Clusters.
S. Maeda, Y. Watanabe, and K. Ohno, *J. Chem. Phys.* 128, 144111-(1,11) (2008).
- 22) DFT Study on Isomerization and Decomposition of Cuprous Dialkyldithiophosphate and Its Reaction with Alkyl Radical.
Y. Luo, S. Maeda, and K. Ohno, *J. Phys. Chem. A* 112(25), 5720-5726 (2008).
- 23) A New Global Reaction Route Map on the Potential Energy Surface of H_2CO with Unrestricted Level.
S. Maeda and K. Ohno, *Chem. Phys. Lett.* 460, 55-58 (2008).
- 24) Intramolecular Vibrational Frequencies of Water Clusters $(H_2O)_n$ ($n=2-5$): Anharmonic Analyses Using Potential Functions based on the Scaled Hypersphere Search Method.
Y. Watanabe, S. Maeda, and K. Ohno, *J. Chem. Phys.* 129, 074315-(1,9) (2008).
- 25) Automated Exploration of Reaction Channels.
K. Ohno and S. Maeda, *Physica Scripta* 78, 058122 (8pp) (2008).
- 26) Decomposition of Alkyl Hydroperoxide by a Copper (I) Complex: Insights from Density Functional Theory.
Y. Luo, S. Maeda, and K. Ohno, *Tetrahedron Letters* 49, 6841-6845 (2008).
- 27) Lowest Transition State for the Chirality-Determining Step in $Ru\{(R)\text{-BINAP}\}$ -Catalyzed Asymmetric Hydrogenation of Methyl-3-Oxobutanoate.
S. Maeda and K. Ohno, *J. Am. Chem. Soc.* 130(51), 17228-17229 (2008).
- 28) Water-Catalyzed Gas-Phase Reaction of Formic Acid with Hydroxyl Radical: A Computational Investigation.
Y. Luo, S. Maeda, and K. Ohno, *Chem. Phys. Lett.* 469(1-3), 57-61 (2009).
- 29) Automated Exploration of Stable Isomers of $H^+(H_2O)_n$ ($n=5-7$) via Ab Initio Calculations: An Application of the Anharmonic Downward Distortion Following Algorithm.

- Y. Luo, S. Maeda, and K. Ohno, *J. Comp. Chem.* 30(6), 952-961 (2009).
- 30) Automated Global Mapping of Minimum Energy Points on Seams of Crossing by the Anharmonic Downward Distortion Following Method: A Case Study of H₂CO.
S. Maeda, K. Ohno and K. Morokuma, *J. Phys. Chem. A* 113(8), 1704-1710 (2009).
- 31) Systematic Search for Isomerization Pathways of Hexasilabenzene for Finding its Kinetic Stability.
M. Moteki, S. Maeda, and K. Ohno, *Organometallics* 28(7), 2218-2224 (2009).
- 32) An Automated and Systematic Transition-Structure Explorer in Large Flexible Molecular Systems Based on Combined Global Reaction Route Mapping and Microiteration Methods.
S. Maeda K. Ohno, and K. Morokuma, *J. Chem. Theory Comput.* 5, 2734-2743 (2009).
- 33) A Systematic Study on the RuHCl-BINAP Catalyzed Asymmetric Hydrogenation Mechanism by the Global Reaction Route Mapping Method.
Koichi Ohno and Satoshi Maeda, *J. Mol. Cat. A Chemical* 324, 133-140 (2010).
- 34) Synthesis and Structures of Stable Base-Free Dialkylsilanimines.
T. Iwamoto, N. Ohnishi, Z. Gui, S. Ishida, H. Isobe, S. Maeda, K. Ohno, and M. Kira, *New. J. Chem.* 34, 1637-1645 (2010).
- 35) Updated Branching Plane for Finding Conical Intersections without Coupling Derivative Vectors.
S. Maeda, K. Ohno, and K. Morokuma, *J. Chem. Theory Comput.* 6, 1538-1545 (2010).
- 36) A Theoretical Study on the Photodissociation of Acetone: Insight into the Slow Intersystem Crossing and Exploration of Nonadiabatic Pathways to the Ground State.
S. Maeda, K. Ohno, and K. Morokuma, *J. Phys. Chem. Letters* 1, 1841-1845 (2010).
- 37) Theoretical Investigation of the Reaction Pathway of O Atom on Si(001)-(2x1).
Shin-ya Ohno, Ken-ichi Shudo, Masatoshi Tanaka, Satoshi Maeda, and Koichi Ohno, *J. Phys. Chem. C* 114(37), 15671-15677 (2010).
- 38) Long-Range Migration of a Water Molecule to Catalyze a Tautomerization in Photoionization of the Hydrated Formamide Cluster.
S. Maeda, Y. Matsuda, S. Mizutani, A. Fujii, and K. Ohno, *J. Phys. Chem. A* 114, 11896-11899 (2010).
- 39) Ab initio anharmonic calculations of vibrational frequencies of benzene by means of efficient construction of potential energy functions.
K. Ohno and S. Maeda, *Chem.Phys.Lett.* 503(4-6), 322-326 (2011).
- 40) Automated Exploration of Chemical Reaction Pathways,
Koichi Ohno and Satoshi Maeda, *Mol. Sci.* 5, A0042 (2011).
- 41) Systematic Exploration of Chemical Structures and Reaction Pathways on the Quantum Chemical Potential Energy Surface by Means of the Anharmonic Downward Distortion Following Method,
Koichi Ohno and Yuto Osada, "*Advances in the Theory of Quantum Systems in Chemistry and Physics*", Springer, in press.
- 42) Exploring Multiple Potential Energy Surfaces: Photochemistry of Small Carbonyl Compounds.
Satoshi Maeda, Koichi Ohno, and Keiji Morokuma, "*Advances in Physical Chemistry*", Vol. 2012, 268124 (13 pages) (2012).